

## Referenser till Vetenskaplig kommentar om värdet av symtomskattning i livets slut

1. Slotman E, Hedman C, Fransen HP, van der Linden YM, Raijmakers NJH, Lundström S. Pain at the end of life in patients with cancer: a population-based study on prevalence, relief, and the role of pain assessment. *Support Care Cancer*. 2026;34(2):116.
2. Fallon M, Walker J, Colvin L, et al. Pain management in cancer center inpatients: a cluster randomized trial to evaluate a systematic integrated approach-the Edinburgh pain assessment and management tool. *J Clin Oncol*. 2018; 36(13):1284–1290
3. Basch E, Schrag D, Henson S et al. Effect of electronic symptom monitoring on patient-reported outcomes among patients with metastatic cancer: a randomized clinical trial. *JAMA*, 2022, 327(24):2413–2422 12.
4. Basch E, Deal AM, Kris MG et al. Symptom monitoring with patient-reported outcomes during routine cancer treatment: a randomized controlled trial. *J Clin Oncol*, 2015, 34(6):557–565. [https:// doi.org/10.1200/JCO.2015.63.0830](https://doi.org/10.1200/JCO.2015.63.0830)
5. Tegenborg S, Fransson P, Martinsson L. The Abbey Pain Scale: not sufficiently valid or reliable for assessing pain in patients with advanced cancer. *Acta Oncologica*. 2023, 62(8), 953–960.